

Cactus Trees, and Lower Bounds on the Spectral Radius of Vertex-Transitive Graphs

Laurent Bartholdi

Abstract. This paper gives lower bounds on the spectral radius of vertex-transitive graphs, based on the number of “prime cycles” at a vertex. The bounds are obtained by constructing circuits in the graph that resemble “cactus trees”, and enumerating them. Counting these circuits gives a coefficient-wise underestimation of the Green function of the graph, and hence an underestimation of its spectral radius.

The bounds obtained are very good for the Cayley graph of surface groups of genus $g \geq 2$, with standard generators (these graphs are the 1-skeletons of tessellations of hyperbolic plane by $4g$ -gons, $4g$ per vertex). We have for example for $g = 2$

$$0.662418 \leq \|M\| \leq 0.662816,$$

and for $g = 3$

$$0.552773 \leq \|M\| \leq 0.552792.$$

1. Introduction: Groups

Throughout this paper, Γ will be a group generated by a finite, symmetric set S , of cardinality $\#S = d$. “Symmetric” means that $S = S^{-1}$. Many of the objects we define will depend heavily on the choice of S , even though we will not make it explicit in the notation.

The *Cayley graph* $\mathfrak{G}$ of Γ is the graph with vertex set Γ , and vertices connected under the right action of S ; i.e. γ and γs are joined for all $\gamma \in \Gamma$ and $s \in S$. The *Markov operator* $M : \ell^2(\Gamma) \rightarrow \ell^2(\Gamma)$ is defined by

$$(Mf)(\gamma) = \frac{1}{d} \sum_{s \in S} f(\gamma s).$$

It is used to study the simple random walk on Γ ; for instance, the probability of return in n steps is $p_n = \langle M^n \delta_1 | \delta_1 \rangle$, where δ_1 is the Dirac function at $1 \in \Gamma$.

The first step in understanding M is the computation of its (operator) norm $\|M\|$, also called the *spectral radius* of $\mathfrak{G}$; indeed, the probabilities p_n satisfy $\limsup_{n \rightarrow \infty} \sqrt[n]{p_n} = \|M\|$. Harry Kesten showed in [Kes59b, Kes59a] the group-theoretical importance of $\|M\|$: we always have

$$\frac{2\sqrt{d-1}}{d} \leq \|M\| \leq 1, \quad (1)$$

with equality on the left if and only if $\mathfrak{G}$ is a tree, i.e. Γ is a free product of $\mathbb{Z}/2$'s and $\mathbb{Z}$'s, with S consisting of the standard generators and their inverses; and equality holds on the right if and only if Γ is amenable.

Assume now that Γ is not free; say it has a relation of length $k \geq 3$. William Paschke obtained in [Pas93] the estimate

$$\|M_\Gamma\| \geq \min_{s>0} \left\{ 2 \cosh(s) + (d-2)Q \left(\frac{\cosh(ks) + 1}{\sinh(s) \sinh(ks)} \right) \right\}, \quad (2)$$

where $Q(t) = (\sqrt{t^2 + 1} - 1)/t$.

The purpose of this paper is to show that the lower bound (2) on $\|M\|$ can be improved if additional hypotheses are made on the number of relations of Γ , and on the number of distinct cyclic permutations of these relations.

Definition 1.1. The *Green function* of Γ is the formal power series

$$G(t) = \sum_{w \in S^* : w \equiv_\Gamma 1} t^{|w|} = \sum_{n \geq 0} p_n d^n t^n,$$

i.e. the growth series of the words representing 1 in Γ .

For two power series $G(t) = \sum_{n \geq 0} g_n t^n$ and $H(t) = \sum_{n \geq 0} h_n t^n$, define $G \lesssim H$ to mean $g_n \leq h_n$ for all $n \geq 0$.

A *prime relator* is a word $w \in S^*$ such that $w \equiv_\Gamma 1$, and such that $v \not\equiv_\Gamma 1$ for all proper subwords v of w .

A set $\mathcal{R}$ of prime relators satisfies the *small cancellation condition* $\mathcal{O}(\eta)$ if for any $w, w' \in \mathcal{R}$, and any factorization $w = uv$ and $w' = vu'$ we have either $u = u'$ or $|v| \leq \eta \cdot \min\{|w|, |w'|\}$.

Note that a prime relator is necessarily a cyclically freely reduced word. The symbol “ $\mathcal{O}$ ” stands for “overlap”. It is a notion close, but strictly weaker than the $C(\eta)$ in small cancellation theory.

The main result of this paper, stated for finitely generated groups, is the following. See in Subsection 2.1 the more general form stated for vertex-transitive graphs:

Corollary 1.2. *Let Γ be a group generated by a finite symmetric set S of cardinality d , and let M be its Markov operator. Assume that Γ has a set $\mathcal{R}$ of prime relators satisfying a small cancellation condition $\mathcal{O}(\eta)$.*

Let $f(t) = \sum_{w \in \mathcal{R}} t^{|w|}$ be the growth series of $\mathcal{R}$, and let ζ satisfy $\zeta - 1 + (1 - 1/d)\zeta f(\zeta^{\eta-1}/(d-1)) = 0$. Construct the power series

$$\begin{aligned} h(t) &= \frac{2(d-1)}{d-2+d\sqrt{1-4(d-1)t^2}}, \\ H(t, u) &= \frac{1-(1-u)^2t^2}{1+(1-u)(d-1+u)t^2} h\left(\frac{t}{1+(1-u)(d-1+u)t^2}\right), \\ g_1(t, u) &= H(\zeta t, u), \\ g_2(t) &= g_1\left(t \frac{d-f(t)}{d-(d-1)f(t)}, \frac{(d-2)f(t)}{d-f(t)}\right), \\ g_3(t) &= h(t)g_2\left(\frac{1-\sqrt{1-4(d-1)t^2}}{2(d-1)t}\right). \end{aligned} \quad (*)$$

Then the Green function of $\mathfrak{G}$ satisfies $G \gtrsim g_3$.

Let ρ be the radius of convergence of g_3 . Then $\|M\| \geq 1/(d\rho)$.

Note that Corollary 1.2 does not supersede Paschke's result (2), in that the bound it gives for the group $(\mathbb{Z}/k) * (\mathbb{Z}/2) * \cdots * (\mathbb{Z}/2)$ is inferior to Paschke's. It does, however, give a superior bound for many groups, and in particular surface groups.

1.1. Surface groups

Consider the fundamental group of a surface of genus $g \geq 2$

$$\Gamma_g = \langle a_1, b_1, \dots, a_g, b_g \mid [a_1, b_1] \dots [a_g, b_g] = 1 \rangle.$$

This group is non-elementary hyperbolic, hence non-amenable; its spectral radius is $O(g^{-1/2})$ for large g . Simple lower and upper bounds come respectively from Γ_g being a quotient of a free group of rank $2g$, and containing by Magnus' Freiheitssatz a $(2g-1)$ -generated free subgroup $\langle a_1, b_1, \dots, a_{g-1}, b_{g-1}, a_g \rangle$:

$$\frac{\sqrt{4g-1}}{2g} \leq \|M_g\| \leq \frac{\sqrt{4g-3}+2}{2g};$$

For more details, see [BCCH97]. Many improvements on these bounds were obtained, see [Pas93, CV96, BCCH97, Žuk97, Nag97, BC02]. Currently, the the best known bounds are

Theorem 1.3. *The spectral radius of the surface groups of genus 2 and 3 satisfy*

$$\begin{aligned} 0.662418 &\leq \|M_2\| \leq 0.662816, \\ 0.552773 &\leq \|M_3\| \leq 0.552792. \end{aligned}$$

These upper bounds are due to Tatiana Nagnibeda [Nag97].

1.2. Reduction to graphs

Equations (1) and (2) hold more generally for any vertex-transitive graph, i.e. for any graph whose group of automorphisms acts transitively on vertices. Remember from [Pas93] that there are vertex-transitive graphs with no simply vertex-transitive automorphism group — for instance, the 1-skeleton of a dodecahedron. Such a graph cannot be the Cayley graph of a group.

In the context of d -regular vertex-transitive graphs, $p_n d^n$ is the number of closed paths of length n at any fixed vertex in $\mathfrak{G}$, and $d\|M\|$ is the asymptotic exponential growth rate of these numbers of closed paths.

In just the same way as any group is a quotient of a free group, any d -regular graph is covered by the d -regular tree. Since closed paths in the tree remain closed in the quotient graph, the spectral radius of any d -regular graph is bounded from below by the spectral radius of the d -regular tree; this proves the left inequality of (1).

Similarly, any vertex-transitive, d -regular graph with a loop of length k at each vertex is covered by the graph $P_{k,d}$ obtained from a $k(d-2)$ -regular tree by replacing each vertex by a k -gon and equidistributing the edges on the k -gon's vertices [Pas93, Proposition 2.4]. The spectral radius of $P_{k,d}$ can be computed using

Theorem 1.4 ([Woe86]; [CS86]; [Bar99, Theorem 9.2]). *Let $\mathcal{X}_1, \mathcal{X}_2$ be vertex-transitive graphs, and let $\mathcal{X}$ be their free product. Let $G_1(t), G_2(t)$ and $G(t)$ be their corresponding Green functions. Then*

$$\frac{1}{(tG(t))^{-1}} = \frac{1}{(tG_1(t))^{-1}} + \frac{1}{(tG_2(t))^{-1}} - \frac{1}{t},$$

where $F^{-1}(t)$ denotes the formal inverse, i.e. the series $E(t)$ such that $E(F(t)) = F(E(t)) = t$.

Indeed if we take $\mathcal{X}_1$ a $d-2$ -regular tree and $\mathcal{X}_d$ a k -cycle, then $\mathcal{X}$ is $P_{k,d}$.

Clearly if $F(t)$ is an algebraic series, then $F^{-1}(t)$ is algebraic of same degree, because $P(t, F(t)) = 0$ implies $P(F^{-1}(t), t) = 0$; also, $\deg(F_1(t) + F_2(t)) \leq \deg F_1(t) \deg F_2(t)$. Simple computations show that for $\mathcal{X}_1$ and $\mathcal{X}_2$ as above G_1 is algebraic of degree 2 and G_2 is algebraic of degree $(k+1)/2$. It follows that $\|M_P\|$ is an algebraic number of degree at most $k+1$.

Another computations shows that $\|M_P\|$ is the right-hand side of (2); the inequality then follows.

Note that Paschke's estimate is valid for all vertex-transitive graphs; the inequality is obtained by constructing a cover for all graphs containing a k -cycle. We note *a posteriori* that $P_{k,d}$ is the Cayley graph of $(\mathbb{Z}/k) * (\mathbb{Z}/2) * \cdots * (\mathbb{Z}/2)$, with $d-2$ copies of $\mathbb{Z}/2$, but there is no *a priori* reason for the graph of smallest norm, $P_{k,d}$, to be the Cayley graph of a group. Our method can also be understood as constructing a transitive graph of minimal norm satisfying some conditions on its cycles; however, this graph of minimal norm will not be the Cayley graph of a group.

2. Preliminaries: Graphs

A *graph* $\mathfrak{G}$ is a pair V, E of sets called respectively *vertices* and *edges*, with maps $\alpha, \omega : E \rightarrow V$ called respectively *start* and *end*, and an involution $\bar{\cdot} : E \rightarrow E$ such that $\alpha(\bar{e}) = \omega(e)$. The *degree* of a vertex is $\deg(v) = \#\{e \in E : \alpha(e) = v\}$. The graph is *d-regular* if each vertex has degree d .

A *path* is a sequence $p = (p_1, \dots, p_n)$ of edges, with $\omega(p_i) = \alpha(p_{i+1})$ for all $i \in \{1, \dots, n-1\}$. Its *length* is $|p| = n$, and its *start* and *end* are $\alpha(p) = \alpha(p_1)$ and $\omega(p) = \omega(p_n)$. It is a *circuit* if $\alpha(p) = \omega(p)$. Paths are multiplied by concatenation; therefore in this definition a graph is nothing but a small $*$ -category with object set V and arrow set E .

A *spike* in a path p is an index $i \in \{1, \dots, |p| - 1\}$ such that $p_i = \overline{p_{i+1}}$. A path is *reduced* if it has no spike. The *spike count* of a path p is the number $\text{sc}(p)$ of spikes in p . A circuit is *prime* if it is reduced, non-trivial, and $\alpha(p_i) \neq \alpha(p_j)$ for all $i \neq j$.

The *Markov operator* on $\mathfrak{G}$ is the operator $M : \ell^2(V, \deg) \rightarrow \ell^2(V, \deg)$ given by

$$(Mf)(v) = \frac{1}{\deg(v)} \sum_{\{e \in E : \alpha(e) = v\}} f(\omega(e)).$$

It governs the behaviour of the simple random walk on $\mathfrak{G}$: given two vertices $v, w \in V$ the probability of going from v to w in n steps is $\langle M^n \delta_w | \delta_v \rangle$, where δ_x is the Dirac function at $x \in V$.

The graph $\mathfrak{G}$ is *transitive* if for any two vertices $v, w \in V$ there is a graph automorphism mapping v to w . Transitivity implies that the probability of return $p_n = \langle M^n \delta_v | \delta_v \rangle$ does not depend on v ; its exponential rate of decay is $\|M\|$, the spectral radius of the random walk.

If Γ is a group with symmetric generating set S , its *Cayley graph* $\mathfrak{G}$ has vertex set Γ and edge set $\Gamma \times S$, with $\alpha(\gamma, s) = \gamma$ and $\omega(\gamma, s) = \gamma s$ and $(\gamma, s) = (\gamma s, s^{-1})$. Clearly, $\mathfrak{G}$ is a transitive graph under the left action of Γ .

Let $\mathfrak{G}$ be a transitive graph, and let $\mathcal{P}$ be a set of prime circuits in $\mathfrak{G}$ at a given vertex v . By transitivity, there exists a translate $\mathcal{P}_w$ of $\mathcal{P}$ at any vertex w . A *spread* of $\mathcal{P}$ is the union of the $\mathcal{P}_w$ for all $w \in V$.

The set $\mathcal{P}$ satisfies the *small cancellation condition* $\mathcal{O}(\eta)$ if it has a spread $\widehat{\mathcal{P}}$ such that for any $p, p' \in \widehat{\mathcal{P}}$, and any factorization $p = qr$ and $p' = rq'$ we have either $q = q'$ or $|r| \leq \eta \cdot \min\{|q|, |q'|\}$.

The *growth series* of a set $\mathcal{P}$ of paths is the formal power series $f(t) = \sum_{p \in \mathcal{P}} t^{|p|}$. The *Green function* of $\mathfrak{G}$ is the growth series of the set of circuits at an irrelevant but fixed vertex.

2.1. Main result

The definitions given above were tailor-cut to make Corollary 1.2 a direct consequence of the following result on graphs:

Theorem 2.1. *Let $\mathfrak{G}$ be a vertex-transitive, d -regular graph, and let $\mathcal{P}$ be a set of prime circuits at a fixed vertex in $\mathfrak{G}$ satisfying $\mathcal{O}(\eta)$. Let $f(t)$ be the growth series of $\mathcal{P}$. Construct the power series $g_1(t), \dots, g_3(t)$ as in (*), Corollary 1.2. Then the Green function of $\mathfrak{G}$ satisfies $G \lesssim g_3$.*


Let ρ be the radius of convergence of g_3 . Then $\|M\| \geq 1/(d\rho)$.

The main idea behind Theorem 2.1 is the construction of *cactus trees* in $\mathfrak{G}$. A *cactus tree* is a circuit in $\mathfrak{G}$ that is built in three stages: at the first stage, a “trunk” is constructed, i.e. a circuit that freely reduces to the trivial circuit. This trunk may not contain any subword of length $(1 - \eta)|p|$ of any prime circuit p .

At the second stage, “fruits”, i.e. prime circuits, are inserted at all vertices of the circuit, in such a way that the resulting circuit is reduced.

At the third stage, “spikes”, i.e. circuits that freely reduce to the trivial circuit, are inserted at all vertices of the cactus tree.

Here is an example of cactus, with the three stages of construction indicated in solid bold, dashed medium and solid thin lines:



Consider the graph $X_{d,m}$ that is the 1-skeleton of the tessellation of the hyperbolic plane by m -gons, d per vertex. Numerically, the growth of cactus trees is a very good underestimation of the growth of circuits in the graph $X_{d,m}$. The Cayley graph of the surface group Γ_g is $X_{4g,4g}$. This explains the bounds given in Subsection 1.1.

3. A Formula: Cogrowth

This section recalls the main result of [Bar99]. Fix a vertex $v \in V$, and consider a set $\mathcal{C}$ of reduced circuits at v . The *saturation* $\langle \mathcal{C} \rangle$ of $\mathcal{C}$ is the closure of $\mathcal{C}$ under the iterated operation of inserting spikes in paths; i.e., $\langle \mathcal{C} \rangle$ is the smallest set of circuits containing $\mathcal{C}$ and such that for all product of paths $pq \in \langle \mathcal{C} \rangle$ and for all $e \in E$ with $\alpha(e) = \omega(p) = \alpha(q)$ we have $p\bar{e}q \in \langle \mathcal{C} \rangle$. (We allow p, q to be empty, in which case by convention $\omega(p) = \alpha(q) = v$).

Let $\mathcal{P}$ be a set of paths. Its *spiky growth series* is the formal 2-variable power series

$$G_{\mathcal{P}}(t, u) = \sum_{p \in \mathcal{P}} t^{|p|} u^{\text{sc}(p)}.$$

The specialization $G_{\mathcal{P}}(t, 1)$ “forgets” the number of spikes in the paths while “remembering” only their lengths, and in case $\mathcal{P}$ consists of all the circuits at v is often called the *Green function* of $\mathfrak{G}$ at v . The specialization $G_{\mathcal{P}}(t, 0)$ counts only reduced paths in $\mathcal{P}$.

Theorem 3.1 ([Bar99], Corollary 2.6). *Let $\mathcal{C}$ be an arbitrary set of reduced paths in a d -regular graph $\mathfrak{G}$, and let $\langle \mathcal{C} \rangle$ be its saturation. Then*

$$\frac{G_{\langle \mathcal{C} \rangle}(t, u)}{1 - (1 - u)^2 t^2} = \frac{G_{\langle \mathcal{C} \rangle}\left(\frac{t}{1 + (1 - u)(d - 1 + u)t^2}, 1\right)}{1 + (1 - u)(d - 1 + u)t^2}.$$

In particular, we have

$$\frac{G_{\mathcal{C}}(t, 0)}{1 - t^2} = \frac{G_{\langle \mathcal{C} \rangle}\left(\frac{t}{1 + (d - 1)t^2}, 1\right)}{1 + (d - 1)t^2}.$$

Assume $\mathfrak{G}$ is transitive and d -regular, and recall p_n denotes the probability of return in n steps of the simple random walk on $\mathfrak{G}$. Then at each vertex there are $d^n p_n$ circuits of length n , and we have the following connection between spectral radius and counting of paths:

Proposition 3.2. *Let $\mathcal{C}$ be a set of circuits at $v \in V$ in a d -regular graph $\mathfrak{G}$, and let ρ be the convergence radius of $G_{\langle \mathcal{C} \rangle}(t, 1)$. Then $\|M\| \geq 1/(d\rho)$.*

3.1. Forbidden words

Let F be a set of words over an alphabet A , and consider the problem of estimating the number of words over A not containing any element of F as a subword. We have the following lemma, which is in essence the Lovasz local lemma:

Lemma 3.3. *Let W be either A^* or the set of reduced words $A^* \setminus \{A^*aa^{-1}A^*\}_{a \in A}$, if A has an involution $a \leftrightarrow a^{-1}$.*

Let $F \subset W$ be a set of “forbidden” words, and set $L = W \setminus WFW$ be the language of words in W not containing an element of F as a subword. Let Φ denote the growth series of F , and let λ, ρ denote the growth rate of L, W respectively; ρ is either $\#A$ or $\#A - 1$.

Then we have $\lambda \geq \rho\zeta$, where ζ satisfies the equation $\zeta - 1 + \frac{\rho\zeta}{\#A}\Phi(\rho^{-1}\zeta^{-1}) = 0$.

Proof. Put on the $S_n = W \cap A^n$ the uniform distribution, and define events Q_j, R_j on S_n as follows: Q_j is the set of words $w \in S_n$ for which there are no factorizations $w = efg$ with $|ef| = j$ and $f \in F$; in other words, Q_j is the event of not containing a forbidden word ending at index j . Set $R_j = Q_1 \cap Q_2 \cap \dots \cap Q_{j-1}$. Note $\mathbb{P}(R_{n+1}) = \mathbb{P}(Q_n|R_n)\mathbb{P}(R_n)$.

We claim that $\mathbb{P}(Q_j|R_j) \geq \zeta$ for all $j \in \{1, \dots, n\}$, and proceed by induction, the basis being $\mathbb{P}(Q_0|R_0) = 1 \geq \zeta$ by definition of ζ :

$$\begin{aligned} \mathbb{P}(Q_j|R_j) &\geq 1 - \sum_{f \in F} \frac{\mathbb{P}(\text{“occurrence of } f \text{ ending at index } j\text{”} \cap R_j)}{\mathbb{P}(R_j)} \\ &\geq 1 - \sum_{f \in F} \frac{\mathbb{P}(\text{“occurrence of } f \text{ ending at index } j\text{”} \cap R_{j-|f|+1})}{\mathbb{P}(Q_{j-1}|R_{j-1}) \dots \mathbb{P}(Q_{j-|f|+1}|R_{j-|f|+1})\mathbb{P}(R_{j-|f|+1})} \\ &\geq 1 - \sum_{f \in F} \frac{\#A^{-1}\rho^{1-|f|} \mathbb{P}(R_{j-|f|+1})}{\zeta^{|f|-1}\mathbb{P}(R_{j-|f|+1})} \\ &= 1 - \rho\zeta\Phi(\rho^{-1}\zeta^{-1})/\#A = \zeta. \end{aligned}$$

(In the third equality, we use $\mathbb{P}(\text{“occurrence of } f \text{ ending at index } j\text{”}) = \#A^{-1}\rho^{1-|f|}$ and independence with $R_{j-|f|+1}$.)

Therefore $\#(L \cap S_n) = \rho^n \mathbb{P}(R_{n+1}) \geq (\rho\zeta)^n$. $\square$

As an application, consider a d -regular transitive graph $\mathfrak{G}$, and a spread $\mathcal{P}$ of “forbidden paths” in $\mathfrak{G}$. Fix a vertex $*$ in $\mathfrak{G}$, and let T denote the set of circuits at $*$. By Theorem 3.1, the spiky growth series of T is

$$\Theta(t, 1-u) = \frac{2(d-1)(1-u^2t^2)}{(d-2)(1+u(d-u)t^2) + d\sqrt{(1+u(d-u)t^2)^2 - 4(d-1)t^2}}.$$

Lemma 3.4. *If F is a set of freely reduced words, then with the notation of Lemma 3.3 the spiky growth series of $T \cap L$ is coefficient-wise at least $\Theta(\zeta t, u)$.*

Proof. For $n \in \mathbb{N}$ consider $T \cap A^n$; then ζ^n is a lower bound of the probability, for a word of length n chosen with uniform probability, to belong to L . For any

$i < j \in \{1, \dots, n\}$, if there exists $w \in T \cap A^n$ whose subword $w_i \dots w_j$ is reduced, then the restriction $T \cap A^n \rightarrow A^{j-i+1}$ to indices $i \dots j$ yields a uniform reduced word of length $j - i + 1$. It follows that

$$\zeta^n \leq \mathbb{P}(w \in F | w \in A^n) \leq \mathbb{P}(w \in F | w \in T \cap A^n).$$

□

Note that in case a more detailed description of F is available, strengthenings of the above result are possible. With surface groups in mind, we consider the concrete case $A = \{a_1^{\pm 1}, b_1^{\pm 1}, \dots, a_g^{\pm 1}, b_g^{\pm 1}\}$; W either A^* or the set F_A of reduced words over A ; and $r = [a_1, b_1] \dots [a_g, b_g]$. We take for F the set of subwords of the cyclic permutations of $r^{\pm 1}$ of length $4g - 1$. As above we write ρ for the growth rate of W . Let $\Lambda(t)$ denote the growth series of L . Then

$$\Lambda \lesssim \frac{1}{1 - \rho t + 8gt^{4g-1} \frac{1-t}{1-t^{4g-1}}}. \quad (3)$$

Indeed, Lemma 3.3 applies, and can be slightly improved by letting F_i denote the set of subwords of length i of $rrr \dots$ and $r^{-1}r^{-1}r^{-1} \dots$, for all $r \in F$, and writing Φ_i for the generating series of F_i . The equation defining Λ is then

$$\Lambda = \frac{1}{1 - \rho t} \left(1 - (\Phi_{4g-1} - \Phi_{4g} + \Phi_{8g-2} - \Phi_{8g-1} + \dots) \Lambda \right);$$

indeed a word in W is either in L , or is of the form fgh , with $h \in L$ and $g \in F_i$ for i maximal. The result follows by inclusion-exclusion.

4. Main Result: Proof of Theorem 2.1

Assume throughout this section that a vertex $v \in V$ has been fixed in the transitive, d -regular graph $\mathfrak{G}$. By Proposition 3.2, a lower bound on $\|M\|$ can be obtained by evaluating $G_{\mathcal{C}}(t, 1)$ for some set $\mathcal{C}$ of circuits at v .

Let $\mathcal{P}$ be a set of prime circuits at v satisfying the small condition condition $\mathcal{O}(\eta)$, and let $f = G_{\mathcal{P}}(t, 1)$ be its growth series. Let $\widehat{\mathcal{P}}$ be an arbitrary but fixed spread of $\mathcal{P}$, i.e. a choice of a translate of every circuit in $\mathcal{P}$ at every vertex of $\mathfrak{G}$.

Start with the set $\mathcal{C}_0 = \{\cdot\}$ consisting only of the empty circuit; its growth series is $G_{\mathcal{C}_0} = 1$. Let $\widetilde{\mathcal{C}}_1$ be the saturation $\langle \mathcal{C}_0 \rangle$ of $\mathcal{C}_0$, and let $H(t, u)$ be its spiky growth series, obtained via Theorem 3.1.

Let $\mathcal{C}_1$ be the set of circuits $c \in \widetilde{\mathcal{C}}_1$ such that, for all $p \in \widehat{\mathcal{P}}$, the circuit c does not contain any prefix p' of p with $|p'| \geq (1 - \eta)|p|$.

Since $f(t)$ counts the prime circuits p , we get $f(t^{1-\eta})$ as the growth of “forbidden” prefixes p' . Solve $\zeta - 1 + (1 - 1/d)\zeta f(\zeta^{\eta-1}/(d-1)) = 0$ for ζ ; then by Lemma 3.4 the spiky growth series of $\mathcal{C}_1$ is bounded from below as

$$G_{\mathcal{C}_1}(t, u) \gtrsim H(\zeta t, u).$$

Consider next $\mathcal{C}_2$, the circuits obtained from $\mathcal{C}_1$ by inserting at each vertex a non-negative number of prime circuits such that the resulting circuit is reduced. Insertion of 0 prime circuits can be done in 0 or 1 ways, depending on whether the vertex is a spike or not, and insertion of $i \geq 1$ circuits can be done in at least $(1 - \frac{2}{d})(1 - \frac{1}{d})^{i-1} f(t)^i$ ways, counting the insert's length. Indeed, to guarantee that the resulting circuit is reduced, it suffices to forbid one out of d starting edges for all prime circuits inserted, except for the last one, for which a starting and an ending edge must be forbidden.

Summing over i gives the generating functions

$$\frac{(d-2)f(t)}{d - (d-1)f(t)}, \quad \frac{d - f(t)}{d - (d-1)f(t)}$$

counting possible insertions at a spike and non-spike vertex respectively. The growth series of $\mathcal{C}_2$ is therefore minorized as

$$G_{\mathcal{C}_2}(t, 1) \succeq G_{\mathcal{C}_1} \left(t \frac{d - f(t)}{d - (d-1)f(t)}, \frac{(d-2)f(t)}{d - f(t)} \right).$$

Finally let $\mathcal{C}_3 = \langle \mathcal{C}_2 \rangle$; a last application of Theorem 3.1 gives a lower bound for $G_{\mathcal{C}_3}(t, 1)$, which in turn, using Proposition 3.2, gives a lower bound on $\|M\|$.

Proof of Theorem 2.1. The proof relies on the construction of cactus trees described in this section. In a cactus tree, i.e. a path constructed as above, mark the edges with the alphabet $\{1, 2, 3\}$ according to the stage at which that edge appeared in the cactus tree.

The arguments given above show that the growth series $G_{\mathcal{C}_3}(t, 1)$ undercounts marked cactus trees; indeed different choices of initial tree (in $\mathcal{C}_1$), prime circuits (in $\mathcal{C}_2$) or final spikes (in $\mathcal{C}_3$) yield different marked cactus trees. To show that $G_{\mathcal{C}_3}(t, 1)$ undercounts circuits, it therefore suffices to show that the markings on cactus trees are uniquely determined; i.e. that two distinct marked cactus trees remain distinct after the marks are erased.

Consider a cactus tree. After removal of all spikes, it gives rise to a unique reduced path; in other words, the order in which the spikes are removed does not change the resulting reduced path.

Now, in this reduced circuit, locate the first subword in $\widehat{\mathcal{P}}$, and pluck it; and repeat till no such subword can be removed. These subwords are necessarily the prime circuits that were inserted in constructing $\mathcal{C}_2$; the only other possibility would be that some circuit $p = p'p'' \in \mathcal{C}_1$ is such that $p'qp''$ contains a prime circuit r at a position before q . This r would then be either a subword of p' , which is forbidden by our construction of $\mathcal{C}_1$, or a subword of $p'q$ containing a part of q ; by the small cancellation condition, a large part of r would subsist in p' , and this is also forbidden by our construction. $\square$

5. Computations: Surface groups

Here we make explicit the arguments in the previous section. Even though our main motivation is to obtain lower bounds for the spectral radius of surface groups, all computations are performed on $X_{d,m}$, the 1-skeleton of a tessellation of the hyperbolic plane by m -gons, d per vertex, introduced above.

Fix a base vertex v of $X_{d,m}$. Consider as prime circuits the $2d$ distinct m -gons touching v , in both orientations. Then $\widehat{\mathcal{P}}$ is the set of m -gons of $X_{d,m}$, taken with all possible base points and orientations.

This choice amounts to taking $f(t) = 2dt^m$. Since two m -gons touch in at most one edge, we have a $\mathcal{O}(1/m)$ small cancellation condition, i.e. we take $\eta = 1/m$.

By Equation (3), we find ζ by solving $\zeta - 1 + 2(d\zeta - 1)/(d^m\zeta^m - 1) = 0$ for ζ 's largest positive root. Numerically, this root is close to 1, and slightly smaller; i.e. $\zeta = 1 - \mathcal{O}(e^{-d})$. For $d = 8$ and $d = 12$, we obtain respectively

$$\zeta(8) = 1 - 0.63 \cdot 10^{-5}, \quad \zeta(12) = 1 - 0.29 \cdot 10^{-10}.$$

We now follow the steps of Theorem 2.1 for $d = m = 8$, corresponding to the surface group of genus 2. We obtain

$$\begin{aligned} h(t) &= \frac{2 \cdot 7}{6 + 8\sqrt{1 - 4 \cdot 7t^2}}, \\ H(t, u) &= \frac{1 - (1 - u)^2 t^2}{1 + (1 - u)(7 + u)t^2} h\left(\frac{t}{1 + (1 - u)(7 + u)t^2}\right), \\ g_1(t, u) &= H(t\zeta(8), u), \\ g_2(t) &= g_1\left(t \frac{d - f(t)}{d - (d - 1)f(t)}, \frac{(d - 2)f(t)}{d - f(t)}\right) \\ &\cong \frac{14(1 - t^2)(1 - 14t^8)}{(1 + 7t^2 - 14t^8 - 2t^{10}) \left(6 + 8\sqrt{1 - 28 \left(\frac{\zeta(8)t(1 - 2t^8)}{1 + 7t^2 - 14t^8 - 2t^{10}}\right)^2}\right)}, \\ g_3(t) &= h(t)g_2\left(\frac{1 - \sqrt{1 - 28t^2}}{14t}\right). \end{aligned}$$

In particular, g_1 and g_2 are algebraic functions of degree 2, and g_3 is algebraic of degree 4. Clearly g_3 's radius of convergence is given by the $g_2(\cdot)$ term, since g_2 counts more circuits than h ; the radius of convergence of g_2 is given by the vanishing of the surd expression

$$1 - 28 \left(\frac{\zeta(8)t(1 - 2t^8)}{1 + 7t^2 - 14t^8 - 2t^{10}} \right)^2,$$

which is a polynomial equation of degree 20, with least solution $\alpha \approx 0.357936$. Now the radius of convergence of g_4 is the minimal t such that $(1 - \sqrt{1 - 28t^2})/14t = \alpha$, namely $\rho = \alpha/(1 + 7\alpha^2) \approx 0.188702$. We then have $\|M_2\| \geq 1/(8\rho)$, so

$$\|M_2\| \geq 0.662418.$$

Similar computations give

$$\|M_3\| \geq 0.552773.$$

The best upper bounds were obtained by Tatiana Nagnibeda [Nag97], by applying Gabber's lemma [GG81] to a function on the edges of $X_{4g,4g}$ defined by the edge-origin's cone type. She obtained

$$\|M_2\| \leq 0.662816, \quad \|M_3\| \leq 0.552792.$$

5.1. Isoperimetric constants

Let $\mathfrak{G}$ be a connected graph. For a subset $K \subset V$ of vertices denote by $\partial K = \{e \in E : \alpha(e) \in K \text{ and } \omega(e) \notin K\}$ the *boundary* of K . The number

$$\iota(\mathfrak{G}) = \inf_{\text{finite, non-empty } K \subset V} \frac{\#\partial K}{\#K} \quad (4)$$

is called the *(edge-)isoperimetric constant* of $\mathfrak{G}$. It is connected to the spectral radius by the following

Theorem 5.1 ([Moh88, MW89]). *Let $\mathfrak{G}$ be an infinite d -regular graph. Then one has*

$$\frac{d^2 - (d-2)\iota(\mathfrak{G})}{d^2 + \iota(\mathfrak{G})} \leq \|M_{\mathfrak{G}}\| \leq \sqrt{1 - \iota(\mathfrak{G})^2/d^2}. \quad (5)$$

Note that, by (1) and (5), $\mathfrak{G}$ is amenable if and only if $\iota(\mathfrak{G}) = 0$.

The isoperimetric constant $\iota(X_{d,m})$ has recently been computed independently by Yusuke Higuchi and Tomoyuki Shirai [HS00] and by Olle Häggström, Johan Jonasson and Russell Lyons [HJL01]. They obtained the values

$$\iota(X_{d,m}) = (d-2) \sqrt{1 - \frac{4}{(d-2)(m-2)}}. \quad (6)$$

In particular, they obtained $\iota(X_{8,8}) = 4\sqrt{2}$ which, together with (5) gives

$$0.431 \approx \frac{16 - 6\sqrt{2}}{16 + \sqrt{2}} \leq \|M_2\| \leq \frac{1}{\sqrt{2}} \approx 0.707.$$

Evidently, these bounds are much weaker than the ones given in this paper. It may be worthwhile to improve the connection between the isoperimetric constant and the spectral radius — probably Theorem 5.1 is not the last word in this topic.

Acknowledgment. The author thanks Tullio Ceccherini-Silberstein, Pierre de la Harpe, Russ Lyons, Tatiana Nagnibeda, and Yuval Peres, who provided valuable comments and encouraged him to write this note.

Laurent Bartholdi,
 Department of Mathematics.
 University of California,
 94720 Berkeley, USA
 E-mail: laurent@math.berkeley.edu

References

- [Bar99] Laurent Bartholdi, *Counting paths in graphs*, Enseignement Math. **45** (1999), 83–131.
- [BC02] Laurent Bartholdi and Tullio G. Ceccherini-Silberstein, *Growth functions and random walks on surface graphs*, to appear in Monatsh. Math., 2002.
- [BCCH97] Laurent Bartholdi, Serge Cantat, Tullio G. Ceccherini-Silberstein, and Pierre de la Harpe, *Estimates for simple random walks on fundamental groups of surfaces*, Colloq. Math. **72** (1997), no. 1, 173–193.
- [CS86] Donald I. Cartwright and Paolo M. Soardi, *Random walks on free products, quotients and amalgams*, Nagoya Math. J. **102** (1986), 163–180.
- [CV96] Pierre-Alain Cherix and Alain Valette, *On spectra of simple random walks on one-relator groups*, Pacific J. Math. **175** (1996), no. 2, 417–438, With an appendix by Paul Jolissaint.
- [GG81] Ofer Gabber and Zvi Galil, *Explicit constructions of linear-sized superconcentrators*, J. Comput. System Sci. **22** (1981), no. 3, 407–420, Special issued dedicated to Michael Machtey.
- [HJL01] Olle Häggström, Johan Jonasson, and Russell Lyons, *Explicit isoperimetric constants and phase transitions in the random-cluster model*, to appear in Ann. Prob.; available at <http://arxiv.org/math.PR/0008191>, 2001.
- [HS00] Yusuke Higuchi and Tomoyuki Shirai, *Isoperimetric constants of (d, f) -regular planar graphs*, preprint, 2000.
- [Kes59a] Harry Kesten, *Full Banach mean values on countable groups*, Math. Scand. **7** (1959), 146–156.
- [Kes59b] Harry Kesten, *Symmetric random walks on groups*, Trans. Amer. Math. Soc. **92** (1959), 336–354.
- [Moh88] Bojan Mohar, *Isoperimetric inequalities, growth, and the spectrum of graphs*, Linear Algebra Appl. **103** (1988), 119–131.
- [MW89] Bojan Mohar and Wolfgang Woess, *A survey on spectra of infinite graphs*, Bull. London Math. Soc. **21** (1989), no. 3, 209–234.
- [Nag97] Tatiana Nagnibeda, *An upper bound for the spectral radius of a random walk on surface groups*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **240** (1997), no. Teor. Predst. Din. Sist. Komb. i Algoritm. Metody. 2, 154–165, 293–294.
- [Pas93] William L. Paschke, *Lower bound for the norm of a vertex-transitive graph*, Math. Z. **213** (1993), no. 2, 225–239.

- [Woe86] Wolfgang Woess, *Nearest neighbour random walks on free products of discrete groups*, Boll. Un. Mat. Ital. B (6) **5** (1986), no. 3, 961–982.
- [Żuk97] Andrzej Żuk, *A remark on the norm of a random walk on surface groups*, Colloq. Math. **72** (1997), no. 1, 195–206.